

Opioid Use and Employment Outcomes: Evidence from the U.S. Military[†]

Abby Alpert
University of Pennsylvania and NBER

Stephen D. Schwab
U.S. Army & Baylor University

Benjamin Ukert
Texas A&M University

December 2022

Abstract

There is significant interest in understanding the labor market consequences of the opioid epidemic, but little is known about how opioid use affects on-the-job performance. We analyze the impact of opioid initiation on workforce performance outcomes by linking individual-level medical and personnel data for active duty military service members. Exploiting quasi-random assignment of patients to physicians in the emergency department, we find that service members assigned to a high-intensity opioid prescribing physician are subsequently less likely to be promoted and more likely to receive disciplinary actions and leave the military. Our results highlight the important productivity costs of opioid use.

[†] We are grateful for helpful comments from Anne Burton, Mireille Jacobson, Kelli Marquardt, Alex Olssen, David Powell, Jonathan Zhang and seminar and conference participants at the APPAM Fall Research Conference, Conference of the American Society of Health Economists, Lehigh University, and Midwest Health Economics Conference. The views and information presented are those of the authors and do not represent the official position of the Defense Health Agency, or the Departments of Army, Department of Defense, or U.S. Government. Author Emails: alpertab@wharton.upenn.edu; stephen_schwab@baylor.edu; bukert@tamu.edu

1. Introduction

The United States is in the midst of an unprecedented opioid epidemic. Since 1999, opioid prescriptions have quadrupled and there have been almost 500,000 overdose deaths involving opioids (CDC (2021)). While much of the evidence has focused on the mortality and health consequences of this epidemic (Maclean et al. (2021)), the impacts on the U.S. labor market and economy are not well understood. Krueger (2017) conjectured that the massive decline in the labor force participation rate since 2000 could be driven in part by the opioid epidemic, which began around the same time as this decline.¹ Some have also speculated that rising opioid use during the coronavirus pandemic contributed to recent labor shortages (Rockeman (2021)). However, evidence suggests that the direction of causality runs in both directions; while opioid abuse may lead to job loss (e.g., Aliprantis et al. (2019), Harris et al. (2020), Park and Powell (2021)), job loss also spurs more opioid abuse (e.g., Case and Deaton (2015); Hollingsworth et al. (2017), Pierce and Schott (2020), Venkataramani et al. (2020)). Furthermore, a multitude of other factors—such as population aging, trade shocks, and automation—have contributed to the two-decade-long decline in labor supply (Abraham and Kearney (2020)).

Prior research on the labor market consequences of the opioid epidemic has focused on aggregate and extensive margin measures of labor supply—primarily employment. However, 63% of adults who report misusing opioids are currently employed (NSDUH, 2020).² How opioid use impacts workers' performance on the job is largely unknown, yet it could have significant impacts on productivity. In this paper, we provide some of the first evidence on how opioid use affects measures of work capacity, job performance, and productivity using individual-level data. We study how receiving an initial opioid prescription affects an individual's subsequent work performance and the mechanisms leading to job separation.

To do this, we use administrative data from the U.S. military, the largest employer in the country. We link individual-level medical and personnel records for the universe of active duty soldiers from 2008 to 2017. The medical records include medical and prescription drug claims and random drug screenings for active duty military members. Personnel files contain

¹ The CDC dates the first wave of the opioid epidemic as beginning in the 1990s (<https://www.cdc.gov/drugoverdose/epidemic/index.html>).

² <https://www.samhsa.gov/data/report/2019-nsduh-detailed-tables> (see Table 1.60A). This figure represents the proportion of adults ages 18+ reporting misuse of opioids who are employed full-time or part-time.

employment and performance measures, such as promotions and demotions, disciplinary actions, physical fitness metrics, and job separations. The records include detailed codes describing the reason for job separation, allowing us to distinguish between voluntary and involuntary discharges. These detailed measures provide an unusually rich characterization of an individual's performance and workforce capability that goes beyond the aggregate labor supply outcomes previously studied in the literature.

The military setting is ideal for studying the impacts of opioid use on workforce outcomes for several reasons. First, the Military Health System (MHS) provides a data linkage between longitudinal healthcare and personnel workforce records that is not typically available in the U.S. labor market. This allows us to observe how job performance evolves after an initial opioid prescription. Second, we are able to continuously follow a large proportion of members over time due to multi-year enlistment contracts which limit turnover in the military. Third, job performance is frequently evaluated and observed through records of promotions and demotions, and direct performance measures (e.g., physical fitness tests) that are assessed at least annually. These outcomes are rarely observed and collected in the civilian labor market. Fourth, military members are highly exposed to opioids (Peters et al. (2019)) and are at risk for negative long-term consequences of opioid use. Finally, opioid use driven productivity consequences have important policy implications for the military's readiness for future missions.

Our empirical approach leverages the quasi-random assignment of patients to physicians when they visit the emergency department (ED) in the military health system. We show that there is wide variation in opioid prescribing behavior across emergency medicine physicians, even within the same hospital and for patients with the same diagnosis. We use the physician's opioid prescribing propensity as an instrument for whether the patient receives an opioid prescription in the ED. We then estimate the impact of receiving an opioid prescription on long-term opioid use and workforce outcomes. We estimate these effects for opioid-naïve individuals to measure the impact of initial opioid exposure. We focus on the ED because patients do not have the ability to request a specific physician in this setting and opioids are frequently prescribed in the ED—about one-quarter of visits in our sample resulted in an opioid prescription. Prior work has used this strategy to show that receiving an opioid prescription in the ED increases the probability of long-term opioid use, opioid use disorder, and overdose death

(Barnett et al. (2017), Barnett et al. (2019), Eichmeyer and Zhang (2022)). However, this strategy has not been used to study labor market outcomes.

Conditioning on hospital, diagnosis, and other patient characteristics, we find that patients assigned to a physician with a 10 percentage point higher prescribing propensity are 8.6 percentage points more likely to receive an opioid prescription in the ED. In line with the prior literature, our instrumental variable estimates show that receiving an opioid prescription in the ED increases the probability of long-term opioid use (filling more than 180 days supply of opioids in the following year) and doctor shopping—two indicators of potential opioid misuse. We further find that individuals who receive an opioid in the ED are 34% more likely to fail a drug screening test in the year following the visit. However, since the majority of drug test failures are for individuals with a legal opioid prescription, we will show that the screening tests do not alone trigger disciplinary actions or job separations.

While the pain relief benefits of opioids could have positive effects on productivity after an acute injury or condition, we find that opioid initiation has, on average, large downstream negative impacts on workforce measures that reduce the productivity of military personnel. Three main findings lead to this conclusion. First, job performance declines following opioid initiation. Our instrumental variable estimates show that the probability of receiving a job promotion decreases by 5% in the year after obtaining a first opioid prescription (relative to a baseline promotion rate of 28%). Job demotions, which occur more infrequently, increase by about 1%, although this estimate is not statistically significant. These negative job performance effects persist through our two years of follow-up.

Second, we document behavioral problems that contribute to lower productivity and job separations. We find a 12% increase in the likelihood that a military member is flagged for a disciplinary action, i.e., misconduct that could result in discipline or separation, including tardiness, unexcused absences, poor attitude, or not performing assigned duties satisfactorily. In contrast, we do not find evidence of an increase in being flagged for a criminal investigation. Overall, job separations increase by 9% following opioid initiation. Discipline-related separations increase by 26%, explaining almost half of the increase in job separations while the remainder of the increase is due to voluntary separations—non-renewal of contracts and retirement. These voluntary separations could partially reflect changes in preferences for work

due to cognitive and behavioral changes or a response to non-promotion or anticipated disciplinary investigations. We do not find evidence of an increase in being discharged for substance abuse. Military discharges due to substance abuse generally require testing positive for an illegal drug, illicit drug possession, or wrongful use of a drug. However, as we show, most opioid-positive drug tests are accompanied by a legal prescription. Therefore, these drug tests would not lead to a discharge for substance abuse, even though some proportion may be misusing opioids as suggested by our results on doctor shopping and long-term use. These findings demonstrate that while behavioral problems resulting from opioid use lead to discharges, substance abuse is not typically the discharge reason. We also find that receiving an opioid prescription is not related to job separations due to medical disability or death. This is consistent with opioid misuse as the mechanism for the workforce effects and not the medical condition itself, further suggesting that our instrument is uncorrelated with condition severity.

Third, we do not find evidence that a member's physical job performance is limited after opioid initiation. Physical fitness test scores and test passing rates are unchanged following receipt of an opioid prescription. The ability to manage pain may offset the negative effects of opioids in this case. However, an important caveat is that our physical fitness test results exclude the most severely injured who can be excused from testing with a physician's note.

In summary, we find significant negative effects of opioids on work performance and employment, which may impact employers' overall productivity and generate higher recruitment and employee retention costs. These effects are largely driven by behavioral issues rather than limitations to physical work capacity. Moreover, behavioral infractions are more common than criminal misconduct among opioid users. Finally, our heterogeneity analyses show that the likelihood that members transition from opioid initiation to negative employment outcomes is related to socio-demographic characteristics. For example, having less than a college education or pre-existing mental health conditions predicts a greater likelihood of negative workforce outcomes in response to opioid initiation. The findings suggest that policies encouraging safer opioid use and access to substance abuse treatment may enhance productivity.

This study contributes to several lines of research. First, we contribute to the literature on the impact of opioid use on labor market outcomes by using individual-level data to study a rich set of performance and productivity outcomes within the firm. A small, but emerging literature

finds negative effects of opioid use on aggregate employment outcomes using data at the state or county level. Aliprantis et al. (2019) and Harris et al. (2020) find that areas with higher prescription opioid access have lower labor force participation rates.³ Other studies use policy variation from the introduction and reformulation of OxyContin (Park and Powell (2021), Powell (2021), Cho et al. (2021)) and rescheduling of hydrocodone (Beheshti (2022)) to show that labor force participation rates decline following large national or state-level shocks to opioid supply.⁴ However, none of these prior studies provide data on workplace performance or productivity outcomes, so they do not capture the full labor market consequences of opioid use. Furthermore, there is limited evidence using individual-level data due to the difficulty of linking prescriptions with workforce outcomes. Two exceptions are studies using Danish administrative data. Laird and Nielsen (2017) leverage patient movers in Denmark to show that patients moving to high-intensity opioid prescribing physicians have reduced labor force participation and earnings, but no changes in the receipt of sick pay and disability insurance. Also using Danish data, Thingholm (2020) instruments a physician's opioid prescribing propensity with the prescribing propensity of spatially connected peers finding broadly similar results. These studies use variation in opioid exposure at the individual level, similar to our study; however, we add to this literature by introducing quasi-random assignment of physicians to address endogenous physician selection concerns and we also add granular measures of work performance.

Second, we contribute to the nascent research on behavioral issues and criminal activity in the workplace. Long-term opioid use has been associated with a wide variety of cognitive and mental changes including mood alteration, difficulty fulfilling obligations, and less attentiveness that can affect workplace behavior and lead to criminal activity (Winkelman (2018)). However, there is no causal evidence documenting how workplace behavior changes after opioid initiation. Evidence exists on the impact of opioid use on criminal activity, particularly heroin sale and possession, however, the magnitudes of the effects vary widely (Meinhofer (2016), Mallat (2018), Mallat (2020), Dave et al. (2021), Deiana et al. (2021)). Relative to the literature, we find

³ To address the potential for reverse causality, Harris et al. (2020) instrument county-level opioid prescription rates with the concentration of high-volume Medicare prescribers. Aliprantis et al. (2019) study geographic variation in opioid prescription rate growth and instrument for prescriptions using historical state prescribing regulations (i.e., triplicate prescription programs).

⁴ Park and Powell (2021) and Powell (2021) also find negative effects of opioid supply on earnings. While earnings are related to productivity, they are indirect measures of underlying job performance and work capacity.

small effects of opioid use on criminal behavior, but large effects on non-criminal behavioral problems.⁵

Third, we add to the broader literature studying the relationship between medical innovations and labor supply. Access to non-opioid pain medications has been shown to improve labor market outcomes (Garthwaite (2012), Bütikofer and Skira (2018)), although these drugs do not have the same addictive properties as opioids. Generally, prior studies have focused on innovations which improve both health and economic well-being (Currie and Madrian (1999)), however, we show that medical innovations, such as those in pain management, can create a tradeoff between health benefits and risks that can have detrimental labor supply effects.

Finally, we show how opioid use has negatively impacted military productivity for active duty service members. Prior research has focused largely on veterans and on the health consequences of opioid use. Opioid abuse among veterans is a growing concern, as opioid-related mortality among veterans increased by 50% from 2000 to 2016 (Lin et al. 2019). Eichmeyer and Zhang (2022) find that opioid initiation in the ED among veterans increases long-term opioid use by 21%, and opioid overdose mortality by 45% within 3 years.⁶ Cesur et al. (2019) find that combat induced opioid abuse leads to annual healthcare costs of \$1.04 billion.

The remainder of the paper proceeds as follows. In Section 2, we provide institutional background on the Military Health System. Section 3 describes the data and Section 4 discusses our empirical strategy. We present the results in Section 5 and Section 6 concludes.

2. Background on the Military Health System

The setting for this paper is the Military Health System (MHS), a distinct entity from the Veterans Health Administration, which is both a payer and provider of care for active duty military service-members, military retirees, and their families. The MHS is a two part system. It provides care in a “direct-care” system that includes 51 military hospitals on military bases and

⁵ Differences in the accessibility of prescription opioids in the military relative to other settings could potentially explain why we do not find significant increases in crime. Prior studies find the largest effects for heroin-related crimes. These studies examine the effects of reduced opioid supply through policy channels, such as PDMPs or the OxyContin reformulation, which lead to substitution to heroin. If opioid prescriptions are more accessible in the military, then we would not expect as much substitution to heroin, and hence lower heroin-related crime rates.

⁶ Barnett et al. (2019) show that veterans’ opioid initiation in the ED increases long-term opioid use. Eichmeyer and Zhang (2023) find similar results on long-term opioid use for veterans exposed to a high prescribing primary care physician, suggesting that the effects may generalize across healthcare service settings.

over 400 outpatient clinics. The MHS also includes the Tricare insurance benefit that pays for medical services both in the direct-care system and in the civilian market (“purchased care”). Active duty military beneficiaries must enroll in “Tricare Prime,” an HMO plan that has near-zero out-of-pocket costs and requires that members receive most care in Military facilities. Active duty members can obtain primary care in the civilian market, but only if they live more than 1 hour away from the nearest military clinic. Additionally, active duty members must have pre-authorization before seeking care in a civilian emergency room or urgent care clinic unless they are traveling and away from their assigned military base. For this reason, most active duty members will receive care at the military base where they reside.

We focus on military emergency departments in this study. MHS EDs are run in a similar fashion as civilian EDs, with the main difference being that they employ active duty physicians. The emergency care physician assignment to a patient in a military hospital is quasi-random. A patient that enters the ED is first triaged by a nurse and then placed in a queue for the next available ED physician. Hospitals vary in how they determine the next available ED physician — i.e., it could be based on the provider’s assigned bed or first-come first-serve (see Chan (2016) for an example) — but at no time can the patient request a specific physician.⁷

3. Data

We use administrative medical data for the universe of active duty soldiers from the Military Health System Data Repository (MDR) covering 2008-2017. The MDR data include medical claims for inpatient and outpatient services as well as pharmaceutical records. These records include all claims for military members regardless of the site of care, including claims from civilian medical providers. We use data from all military services (Army, Navy, Air Force, Marine Corps) for most of our analyses, although we focus on the Army in some specifications where the outcome measure is unavailable for other services.

We link these medical records with a rich dataset of workforce measures that come from multiple military personnel systems including the Defense Manpower Data Center (DMDC) and

⁷ Obermeyer and Chang (2020) find that physicians in one academic hospital have some discretion in selecting patients in the ED. However, this is not the assignment mechanism used in the military health system. In the MHS, doctors are generally assigned to patients based on room capability and capacity. We provide tests of balance on patient characteristics in Section 5 and do not find evidence of selection on observable characteristics.

Integrated Total Army Personnel Database (ITAPD). These data include information on opioid use as measured by drug testing and workforce outcomes such as promotions, demotions, physical fitness, adverse disciplinary actions, and job separations. Demographic variables include age, race, gender, marital status, education, military rank, job title, and military tenure.

3.1 Sample Construction

We construct a sample of emergency department (ED) visits in hospitals located on U.S. military bases. We allocate prescription opioids filled within 7 days of the ED visit to the ED physician.⁸ As described in more detail in Section 4, we compute a leave-out, residualized opioid prescribing rate for each physician as our instrument for receiving an opioid prescription. We restrict our sample to physicians that treat more than 10 patients in the ED in a year. We also limit our sample to ED visits for patients who are opioid-naïve (i.e., patients who did not fill an opioid prescription within the 6 months prior to the index ED visit (Barnett et al. 2017)).⁹ This allows us to measure outcomes associated with opioid initiation. Our final sample includes 1,447,758 ED visits from 2008-2017.

3.2 Outcome Measures

3.2.1 Opioid Prescriptions and Misuse

We use multiple measures to track opioid use in the year following the index ED visit. We examine the number of opioid prescriptions filled and whether the patient received opioid prescriptions from 7 or more different providers—an indicator of “doctor shopping”.¹⁰ Additionally, we measure long-term opioid use, which is defined as filling more than 180 days supply of opioids within the year following the ED visit, excluding the initial prescription. This

⁸ We use the cutoff of 7 days since it is a common metric used in the literature (e.g., Barnett et al. 2017). There is only a slight difference in the number of prescriptions included in our sample if we use smaller windows. This is because most of the opioid prescriptions filled within 7 days of the ED visit are filled on the same day (72%) or within the first 2 days (91%) of the visit. Although relatively rare, we allow for multiple ED visits for the same person and treat each one as a new index visit. However, we exclude any ED visits occurring within 7 days of the initial visit to allow for accurate attribution of the opioid prescription.

⁹ We do not select our sample based on a specific set of diagnosis codes that would be treated by opioids because there are a wide range of conditions that receive opioid prescriptions in the ED for both appropriate and inappropriate reasons (Alpert et al. 2020).

¹⁰ While prior work often uses 5 or more opioid prescribers in the year to measure doctor shopping (e.g., Buchmueller and Carey (2018)) we use a cutoff of 7 or more opioid prescribers because military members generally have a higher number of visits with different providers compared to other populations (only 0.3% of military service members are categorized as exhibiting doctor shopping behavior).

measure, which has been used in prior work (Barnett et al. 2017), is an indicator of potential opioid dependency or misuse since clinical guidelines recommend a much shorter course of treatment for acute medical conditions (CDC (2020)).

As a complementary measure of opioid misuse, we use data from random drug screening tests. The military randomly tests 10% of service members each month and tests 100% of members once per year. We measure whether the patient failed a drug test within one year following the ED visit by type of drug (opioid, heroin, marijuana, benzodiazepines, and all other drugs). We also observe whether the member had an “excused reason” for test failure (i.e., they had been prescribed an opioid). The drug screening data allow us to capture illicit opioid and other drug use not captured in the prescription data.

3.2.2 Job Performance

First, we study promotions and demotions as an observable measure of job performance. Promotions occur frequently in the military. Promotions for the lowest ranks (Private through Private First Class) are largely automatic based on time in service, but there is some discretion in promotion decisions at the margin. Promotions for mid-level enlisted ranks (Specialist to Staff Sergeant), however, are largely merit based. We exclude promotions for officers because they are rare in our data and made through a centralized decision process.¹¹ We also study demotions, which occur less frequently (about 4% of members are demoted each year). Demotions occur when a service member violates the Uniformed Code of Military Justice (UCMJ) and can include relatively minor infractions such as showing up late for work or disrespecting a more senior non-commissioned officer as well as larger infractions that could result in a court-martial.¹²

Second, we study disciplinary actions, formally known as “Suspension of Favorable Personnel Actions (SFPA) Flags.” These records are available for the Army only. We evaluate

¹¹ For other outcomes studied, we include only a small number of officers because we control for AFQT test scores, which only enlisted members take (unless officers began their career as enlisted members). Thus, the difference in sample size between the promotions analysis and the analysis of other outcomes is small.

¹² We measure promotions by evaluating whether a service member’s rank is higher than the rank at the time of the ED visit within 1 and 2 years after the ED visit. Likewise, we evaluate a demotion if the rank is lower than the rank at the time of the ED visit. In some instances, it is possible that a member is promoted and demoted. In such cases, we only count the first change in rank. If multiple promotions and demotions occur (which happens in only <0.2% of cases) we count these events as both promotions and demotions.

the type of SFPA recorded: discipline flag, criminal investigation flag, and flag for drug or alcohol abuse. Having a flag is fairly common, with 22% of individuals receiving at least one in a given year. The “discipline flag” includes minor infractions, such as being late to work and unexcused absences, but excludes drug and alcohol offenses and all other categories such as security violations and domestic violence. The drug or alcohol abuse flag would be used if the member had a positive drug test (without an excused reason) or possession of an illicit drug or failed to complete alcohol or drug abuse rehabilitation programs (see Army Regulation 600-85 and Army Regulation 635-200). Members would typically not be flagged if they tested positive for opioids but had a legal prescription.

Third, we study the probability of obtaining a security clearance which is an indicator of performance issues. Security clearances are essential for performing the tasks of most military jobs and are needed for career advancement (in our sample, 78% hold a security clearance). Obtaining a security clearance involves an extensive background check. A member could be denied a security clearance for reasons including drug/alcohol involvement, criminal conduct, and financial considerations (e.g., running up debt). Once a member has a security clearance, having it revoked is considered a serious violation that would end most military careers. We focus on two outcomes: whether one was denied a security clearance or had it revoked.

3.2.3 Work Capacity

As a direct measure of work capacity, we use data from physical fitness tests to measure an individual’s physical readiness for the job. The Army Physical Fitness Test (APFT) is administered to each Army service member at least once per year, although most service members will take 2 tests per year. The APFT is designed to test physical strength, endurance, and cardio-respiratory fitness. Individuals who are injured are exempt from taking the test if they have a formal medical provider’s note. We study whether individuals take the APFT, pass the APFT, and their test score (which is standardized to the Z-score).

We also evaluate the physical capabilities of Army members with scores from the Physical Capability Grading System. These scores are derived from the Periodic Health Assessment (PHA) that Army members receive annually. In this assessment, a physician conducts a physical exam to assess a member’s current health status and identifies any medical

conditions. They also conduct a behavioral health screening. We focus on the physical and the behavioral assessment Z-score as our outcomes of interest.

3.2.4 *Employment Separation*

Finally, we study job separations as a consequence of poor performance or work capacity. Approximately 18% of service members in our data leave the military within a year of the ED visit. Job separations occur for voluntary and non-voluntary reasons. Voluntary separations generally occur when the contract expires and the service member makes the choice to not sign a new contract.¹³ Non-voluntary exits occur for disciplinary or medical reasons.

Military separations are classified in two distinct ways in our data. First, a “reason code” is entered for the separation. There are more than one-hundred reason codes which are highly descriptive. The most common reason is “expiration of term of service,” or the end of the contract. Other codes include reasons such as: “pattern of minor disciplinary infractions”, “civil court conviction” and “desertion.” We analyze the universe of discharges and create 7 categories to classify the reason codes as follows: non-renewal of contract or retirement, discipline, failure to meet physical standards, substance abuse, other non-discipline, medical, and uncharacterized.

Second, service member exits are separately given a character of separation code that impacts their military benefits: “Honorable”, “General”, “Other than Honorable,” “Bad Conduct”, and “Dishonorable.” An important note is that while the reason code is not punitive in nature, a “General” discharge, for instance, makes the service member ineligible to reenter the military and precludes use of GI bill education benefits. Any discharge below “Honorable” requires legal justification and is administratively burdensome. This means that a service member may exit with a negative reason code but still receive an honorable discharge. Thus, Honorable discharges can be given for both voluntary (e.g., contract end date, transfer, and retirement) and involuntary (e.g., disciplinary actions) separations if the infractions are considered fairly minor. In our analysis, we categorize separations as either “Honorable” or “Non-Honorable” (which includes all discharges excluding Honorable).

¹³ Senior ranks are not required to sign contracts and can remain in the military until they serve a maximum number of years or request to leave. However, the military maintains “retention control points”. For instance, a Staff Sergeant must either be promoted to Sergeant First Class or leave the military after 20 years of active duty service.

4. Empirical Strategy

We estimate the causal effect of receiving an opioid prescription during an ED visit on long-term opioid use and workforce outcomes. Receiving an opioid prescription is correlated with injury severity or pain and these traits themselves predict negative workforce outcomes. To address this endogeneity issue, we exploit variation in physicians' propensities to prescribe opioids. We instrument the receipt of an opioid prescription in the ED with physician-level residualized (leave-one-out) opioid prescribing propensities. We focus on the ED setting where patients do not have the ability to request a specific physician during their visit, creating quasi-random assignment of patients to physicians. Our identification strategy relies on idiosyncratic differences in patients' probability of receiving an opioid prescription stemming only from differences in physician practice styles.

Following a strategy used in prior work (Eichmeyer and Zhang (2022); Barnett et al. (2017)), we construct an instrument that measures physician opioid prescribing intensity. In the first step, we estimate residuals from the following regression:

$$Prescription_{ijt} = \alpha_{ht} + \sigma_{hw} + \theta_d + X'_{it}\beta + \epsilon_{ijt} \quad (1)$$

where $Prescription_{ijt}$ is a binary indicator that equals 1 if patient i received an opioid prescription within 7 days of their ED visit with physician j in month-year t . We control for hospital-month-year fixed effects α_{ht} and hospital-day of week fixed effects σ_{hw} to account for differences in prescribing rates across hospitals and time.¹⁴ We include diagnosis fixed effects θ_d to account for the possibility that some ED physicians may specialize in treating certain diagnoses or higher severity cases and would have a higher tendency to prescribe opioids.¹⁵ We also control for patient demographic characteristics X_{it} , including indicators for age group, White, female, married, college educated, Armed Forces Qualification Test (AFQT) score, military rank, and military tenure. Additionally, we include Military Service-by-occupation fixed effects to account for differences in leadership and culture across military occupational

¹⁴ Month-year takes on unique values for each month-year combination, while day of week takes on seven values. Time of day is not available in the MHS data.

¹⁵ We define diagnoses by the first 3-digits of the primary ICD-9 or ICD-10 code on the ED claim.

specialties (MOS). After conditioning on these fixed effects, the residual variation in the prescribing rate ϵ_{ijt} represents idiosyncratic factors affecting physician prescribing decisions.

For each patient, we then construct our residualized, leave-out instrument of physician opioid prescribing intensity, $Intensity_{ijy}$, as the mean of the physician j 's residuals across the calendar year y from equation (1), leaving out the residual for patient i (denoted by $-i$):

$$Intensity_{ijy} = \frac{1}{N_{-ijy}} \sum_{i=1}^{N_{-ijy}} \hat{\epsilon}_{-ijy} \quad (2)$$

where N_{-ijy} is the number of ED encounters for physician j in year y , excluding patient i . We leave out patient i to avoid bias from including the error term in both the instrument and outcome variables. This instrument allows us to measure differences in the opioid prescribing rate across physicians within the same hospital who are treating the same diagnosis. We also construct an alternative binary instrument used in some specifications where we define physicians as having a prescribing $Intensity_{ijy}$ in the first versus fourth quartile of the distribution (henceforth, “low intensity” and “high intensity” opioid prescribers).¹⁶

Our first stage relationship is estimated with the following equation:

$$Prescription_{ijt} = \delta Intensity_{ijy} + \alpha_{ht} + \sigma_{hw} + \theta_d + X'_{it}\beta + \epsilon_{ijt} \quad (3)$$

We then estimate our second stage using 2SLS, taking the general form:

$$Y_{ijt} = \gamma \widehat{Prescription}_{ijt} + \alpha_{ht} + \sigma_{hw} + \theta_d + X'_{it}\beta + \epsilon_{ijt} \quad (4)$$

where Y_{ijt} measures long-term opioid use or workforce outcomes in the 1-2 years following the initial ED visit. In both equations, we include the full set of controls from equation (1). Standard errors are clustered by physician. We interpret differences in outcomes between patients assigned to higher or lower intensity opioid prescribing physicians as the effect of a change in the probability of receiving an opioid prescription. Identification relies on the assumption that physician assignment is as good as random (i.e., conditional independence) and that our instrument does not operate through channels other than opioid prescribing (i.e., exclusion

¹⁶ In the 1st versus 4th quartile specification, we compute our $Intensity_{ijy}$ instrument including patient i . This allows for physicians to be consistently defined as either high or low intensity prescribers across all patients within a year. Since this is a binary instrument, whether we include or exclude patient i does not meaningfully impact the results.

restriction). Additionally, we assume monotonicity in opioid prescribing intensity across patients. We provide evidence supporting these assumptions in Section 5.

5. Results

5.1 Descriptive Statistics and Balance Test

Figure 1 shows the distribution of the raw mean opioid prescribing rate and the residualized instrument for each physician-year. There is significant variation in both of these measures across ED physicians. The opioid prescribing rate ranges from 14.1% to 30.1% when comparing the mean rate for the bottom and top quartiles of physicians. After controlling for hospital, time, diagnosis, and patient characteristics, we find that moving from the bottom to top quartile physician increases the mean prescribing rate by 19 percentage points.¹⁷

In Table 1, we provide descriptive statistics for the sample of ED patients (Column 1) and a balance test showing the relationship between patient characteristics and our instrument (Columns 2-7), supporting the conditional independence assumption. The average military ED patient is 26 years old, predominately male (79%), White (65%), married (53%), does not have a college degree (95%), has junior rank (62%), AFQT score of 60, and has a military tenure of 5 years. Relative to the commercially insured and Medicare populations, the military ED patient is younger, more likely to be male, and exhibits a lower likelihood of college education, however, the ED opioid prescribing rate of 22% is quite similar to other settings (e.g., Barnett et al (2017), Alpert et al (2020)). To provide evidence of quasi-random assignment of patients to ED physicians, we regress each patient characteristic on our instrument for physician prescribing intensity. Each cell is from a separate regression. Column 2 shows coefficients on the continuous instrument of prescribing intensity and Column 5 shows coefficients on the binary (top vs. bottom quartile intensity) instrument. If patients are randomly assigned to ED physicians, we would expect that the demographic characteristics would be unrelated to prescribing intensity. Indeed, we do not observe statistically significant differences across patients seen by a higher or lower opioid prescribing physician for any of the demographic characteristics. In an alternative version of the balance test, we regress the physician prescribing intensity instrument on all patient characteristics jointly. These results are displayed in Appendix

¹⁷ The 5th, 25th, 75th and 95th percentiles of the raw opioid prescribing rate distribution are: 0.08, 0.15, 0.28, 0.41. The same percentiles of the residualized distribution are: -0.10, -0.04, 0.03, 0.13.

Table 1. Column 1 shows a regression of a binary indicator for receiving an opioid prescription. As expected, this is correlated with patient characteristics. Column 2 shows the balance test. Consistent with random assignment, patient characteristics do not predict the physician opioid prescribing intensity. Almost all coefficients are close to zero and the joint F-statistic is 0.27. In Column 3, we show another variant of the balance test where we construct the opioid prescribing intensity instrument using only hospital-month-year, hospital-day of week, and diagnosis fixed effects. The results are almost identical.¹⁸

5.2 Opioid Prescription and Misuse Outcomes

The existing literature has shown that assignment to a high intensity physician increases the probability of receiving an opioid prescription in the ED and long-term opioid use for veterans and Medicare beneficiaries (Barnett et al (2017), Eichmeyer and Zhang (2022)). In Table 2, we first replicate these findings using our sample of active-duty military members. In Column 1, we estimate the first stage relationship between the physician prescribing intensity and the probability of receiving an opioid prescription following the ED visit. Panel A shows the results for the continuous prescribing intensity instrument and Panel B shows the results for the binary (top vs. bottom quartile) instrument. Both instruments have a strong association with a patient's probability of receiving an opioid prescription. Panel A shows that patients assigned to a physician with a 10 percentage point higher prescribing intensity are 8.6 percentage points more likely to receive an opioid prescription, while Panel B shows that assignment to a physician in the top quartile of prescribing intensity increases the probability of receiving an opioid prescription by 18.5 percentage points compared to the bottom quartile.

In Columns 2-4 we display the results from the 2SLS regressions for long-term opioid use. Column 2 shows the number of opioid prescriptions filled during the first year of follow-up; Column 3 shows an indicator of doctor shopping (having 7 or more prescribers); and Column 4 shows an indicator for long-term opioid use (180 days supply within one year). The results are similar for both instruments and we focus on the continuous instrument, which uses the full sample, in our discussion. For the continuous instrument, the results show that after receiving an

¹⁸ When we replicate Table 1 using only hospital-month-year, hospital-day of week, and diagnosis fixed effects to construct the instruments, we also find similar coefficients and p-values. The balance test results are not sensitive to including sociodemographic controls in the residualization.

opioid prescription in the ED, an individual has 11% more opioid prescription fills (baseline mean fill rate of 0.36), is 60% more likely to exhibit doctor shopping behavior (baseline mean of 0.30%), and is 62% more likely to have long-term opioid use (baseline mean of 0.29%).¹⁹

Table 3 presents complementary evidence of long-term opioid use from random drug screening tests in the year following the ED visit. Unlike the prescription data, drug screenings capture opioid use from both medical and illicit sources, which prior studies have not been able to examine. Column 1 shows that the probability of failing a drug test increases by 1 percentage point (34% increase) after receiving an opioid prescription. Column 2 shows that the increase in drug test failures is predominantly driven by opioid drug test failures for members who have a legal prescription (0.8 percentage point, or 112%, increase). There is no statistically significant increase in drug test failures from opioids without a prescription (Column 3). In Columns 4-7 we look at other drug test failures that could be substitutes (heroin and marijuana) or complements (benzodiazepines) for prescription opioids, and other drugs.²⁰ We find that the probability of a heroin drug test failure decreases by 0.05 percentage points, potentially stemming from patients' expanded access to prescription opioids following an initial opioid prescription.²¹ On the other hand, we find no changes in failed drug tests for marijuana, benzodiazepines, or all other drugs.

The results in this section are broadly consistent with the prior findings in the literature—individuals who encounter a high intensity prescriber in the ED are significantly more likely to initiate long-term opioid use. However, an important question remains: how does opioid initiation (which leads to long-term opioid use) impact work capacity and job performance? We bring to bear our linked healthcare and workforce data to examine this question.

5.3 Workforce Outcomes

¹⁹ In our sample, about one percent of the opioid-naïve who are prescribed an opioid in the ED will have long-term opioid use (0.29/22).

²⁰Alpert et al. (2018) find evidence of substitution between opioids and heroin after OxyContin became abuse-deterrent and Powell et al. (2018) show substitution from opioids to marijuana following medical marijuana law implementation. Concomitant opioid and benzodiazepine use is common, albeit risky (e.g., Hernandez et al., 2018).

²¹Heroin use is generally hard to capture in drug tests because it metabolizes quickly (Cone et al, 1991). The large reduction could be picking up reductions along both the extensive and intensive margins. Patients may be less likely to use heroin at all, but also may use less heroin due to having access to opioids.

In this section, we examine the effects of opioid initiation on job performance, work capacity, and job separations. We then analyze how these outcomes vary across subgroups defined by socio-economic characteristics and pre-existing mental health conditions.

5.3.1 Job Performance

In Table 4, we examine promotions and demotions among enlisted service members as an observable measure of job performance. Promotions are a high frequency outcome in the military with about 28% of enlisted members receiving a promotion within one year of the ED visit and 39% within two years. Demotions for poor job performance, on the other hand, are quite rare (about 4%). Using the continuous instrument in Panel A, we find that the probability of receiving a promotion decreases by 1.4 percentage points within one year of receiving an opioid in the ED and by 1.2 percentage points within two years. These findings imply 5% and 3% fewer promotions, respectively, relative to the baseline means. Results from the binary high intensity instrument are similar. Not surprisingly, the estimates for demotions are noisier given their infrequent occurrence. The probability of being demoted increases by 1.4% after one year (relative to the baseline mean of 3.5%) and 4% in the second year (baseline mean of 4.7%), although these estimates are not statistically different from zero. The larger effects for demotions in the second year could reflect the longer time it takes for demotions to pass through legal and administrative channels, while promotions are more quickly implemented by superiors.

5.3.2 Mechanisms: Physical Work Capacity

Our results on promotions show a sizeable decline in job performance following opioid initiation. We use our detailed workforce data to explore the potential mechanisms driving these results. We first examine how opioids affect physical performance. Similar to many other physically-demanding civilian jobs such as construction, manufacturing, and mining, it is essential for military members to maintain a high level of physical fitness for performing the core functions of their jobs. Opioids, even when taken as prescribed, have known physiological effects on physical performance, such as slowed breathing and heart rate and delayed reaction times, which reduce endurance and make exercise more difficult (Mayo Clinic (2020)). Moreover, opioids can have negative effects on mental functioning and could lead to depression, which may lead military members to put less effort into their training (Mazereeuw et al. (2018), NIDA (2020), Smith (2021)). On the other hand, opioid use could potentially improve physical

performance by reducing the pain from an acute injury or condition. Whether these conflating effects translate into meaningful reductions in work capacity is unknown. The military is a useful setting for studying this question because members are required to take physical fitness tests that have an explicit passing threshold reflecting the physical demands of the job. Thus, we can observe whether opioid initiation impacts workers' ability to meet these physical demands.

Table 5 displays the effects of opioid use on physical performance outcomes from the Army Physical Fitness Test (APFT) in the year after the ED visit. In Column 1, we do not find strong evidence of a decline in the probability of taking this test after receiving an opioid prescription, even though members could receive permission to delay testing due to the injury or medical condition. The coefficients are small and only marginally statistically significant for the continuous instrument. Conditional on taking the APFT, the probability of passing the test decreases by about 0.7%, although this is not statistically significant. The Z-score standardized physical fitness test score decreases by a statistically insignificant 0.03 standard deviations (SD=44 points on a scale of 300 points). Overall, we find little evidence that opioid initiation affects physical fitness scores.

5.3.3 Mechanisms: Behavioral Problems and Criminal Activity

We also consider the effects of opioid use on workplace behavioral problems. In cases of long-term use, opioids have been associated with a wide variety of mental health changes including mood alteration, difficulty fulfilling obligations, lower attention span and less attentiveness (AJMC Perspectives (2020), Meyer (2019), Richards et al. (2018)). More broadly, opioids can increase absenteeism (CDC, 2019), and opioid use is associated with criminal justice system involvement (Winkelman (2018)). All of these factors could lead to problematic workplace behaviors that limit productivity and performance. However, there is little causal evidence for most of these relationships.²²

In Table 6 we evaluate whether receiving an opioid prescription changes the probability that a disciplinary flag (known as a "Suspension of Favorable Personnel Actions (SFPA) Flag") was recorded. There are 19 distinct groups for which an individual may receive a flag for

²² The best available evidence links prescription opioids and crime, though the literature is mixed (Meinhofer (2016), Mallat (2018) & (2020), Dave et al. (2021), Deiana et al. (2021)), and some have suggested that this link may not be as strong as it is for illicit drugs (Maclean et al. (2021)).

problematic behaviors. We look at three specific flag groups where we would expect to see the largest effects of opioids: a flag for discipline, a flag for criminal investigation, and a flag that combines drug and alcohol abuse.²³ The discipline category includes minor infractions such as being late to work and unexcused absences, but excludes drug and alcohol offenses and all other categories such as security violations, and domestic violence.²⁴

We find evidence of a 1.1 percentage point increase (12% relative to the baseline mean of 8.7%) in the probability of having a discipline-related flag within one year of receiving an opioid prescription which is statistically significant at the 5% level. Estimates in year two are somewhat smaller and not statistically significant.²⁵ Column 2 shows results for flags given for criminal investigation. These effects are small and statistically insignificant. However, they become larger in magnitude in the second year, which may be suggestive that criminal behaviors could materialize in the longer run. We also do not find effects of opioid initiation on drug and alcohol flags (Column 3). This is consistent with our earlier drug test results in Table 3. Despite an increase in long-term opioid use, most of these individuals are obtaining legal prescriptions and thus would not be flagged for drug abuse. This highlights the challenge of detecting problematic opioid use in the workplace through drug screening.

Another outcome which may indicate behavioral problems is the probability of receiving a denial of a security clearance or having a security clearance revoked. Obtaining a security clearance involves an extensive background check of a military member's history including drug and alcohol involvement, criminal conduct, and financial records (e.g., running up debt). 78% of Army members hold a security clearance at baseline. In Appendix Table 2, we find that opioid use after an ED visit leads to a higher probability of receiving a denial of a security clearance by 0.5 percentage points after one year (28% increase), but has no effect on the more serious infraction of having one's security clearance revoked.

These findings are consistent with the overall picture of worsening job performance among those who initiated opioid use. From this set of results, it appears that the negative effects of opioids on performance are largely driven by behavioral issues rather than physical

²³ There is no separate flag or data specifically recording absenteeism or missed days of work.

²⁴ The subgroup categories do not sum to the total SFPA flags since we have excluded specific flags that are rarely observed or would be unlikely to be related to opioid use.

²⁵ The less precise estimates in the second year could potentially reflect that some members receiving disciplinary flags in year 1 would be discharged for disciplinary reasons by year 2, as we will show in the next section.

limitations. Moreover, we find that minor behavioral infractions are more common than criminal misconduct among opioid users.

5.3.4 Job Separations

The lower rate of promotions along with the higher rate of disciplinary actions and security clearance denials may eventually lead to involuntary discharges from the military. Voluntary separations— such as contract non-renewal, transfer to the national guard or military reserve, or retirement— may also occur if opioid use changes an individual’s preferences for work. In some cases, a voluntary separation such as contract non-renewal or transfer could be done to pre-empt a future discipline-related discharge or as a response to non-promotion.

In Column 1 of Table 7, we show that the probability of being discharged from the military for any reason increases by 1.6 percentage points within one year of the ED visit. The magnitude of this effect is substantial and represents an increase in separations of 9% relative to the baseline mean of 18%. We find similar effect sizes two years after opioid initiation.

In Columns 2-8, we categorize discharges based on the military’s separation reason code. We find that the largest proportional increase in discharges is for discipline-related reasons,²⁶ which increased by 26% relative to the baseline mean. The number of discipline discharges remained elevated (16% increase) and statistically significant in the second year after the ED visit. Voluntary discharges, which occur when a member does not renew their contract or retires, increased by 17% within the first year, but the estimates are not statistically significant in the second year. Although we observe a faster rate of increase for discipline-related discharges relative to voluntary discharges, they are a smaller share of overall discharges and account for about 42% of the overall discharge effect. Voluntary discharges due to non-renewal of contracts and retirement comprise the majority of the remaining increase in discharges.

For the other discharge types, the estimates are not statistically distinguishable from zero. Finding no effect for discharges due to failure to meet weight and body fat standards is consistent

²⁶ Discipline discharges are given for the following reasons: civil court conviction, security, court martial, AWOL/Desertion, Good of the Service (discharge in lieu of court martial), misconduct, pattern of minor disciplinary infractions, Commission of a Serious Offense, failure to meet minimum qualifications for retention, unsatisfactory performance, unfitness or unacceptable conduct, discreditable incidents, imprisonment/desertion, failure of course of instruction, failure of selection for promotion, motivational problems (apathy).

with our prior results which showed that opioid use did not impact physical fitness test scores. Additionally, we do not find an increase in discharges due to substance abuse. While this might seem counterintuitive, it is consistent with the drug screening test results showing that the majority of opioid-positive drug tests are accompanied by a legal prescription, and therefore, we would expect few discharges due to opioid misuse to be coded as substance abuse. However, given the significant increase in doctor shopping we observe, it is likely that a proportion of these individuals are misusing opioids. It is difficult for the military to discharge individuals for opioid abuse based on drug screenings because of the presence of legal prescriptions. Instead, it is more likely that opioid-related discharges are coded with a reason related to job performance.

The remaining three categories of discharges in Columns 6-8 serve as placebo tests. “Other non-discipline discharges” contain a dozen other reasons for discharge that should be unrelated to opioid use—for example, early release to attend school, pregnancy or parenthood, and errors made by the military in the enlistment process (e.g., underage enlistment). Indeed, we do not find any impact of receiving an opioid on these outcomes. We also examine medical-related discharges which include disability and death. The effects of receiving an opioid prescription on medical discharges are close to zero and statistically insignificant. This further supports our identifying assumption that patients’ assignment to ED doctors is unrelated to the severity of their injury or health condition.

Finally, in Columns 9-10 we examine discharges using the broader “Character of Separation” codes which determine veteran benefits. We report discharges as falling into two separate categories: “Honorable” or “Non-Honorable.” Most service members receive honorable discharges, meaning their service was not marred by anything negative. Given the lengthy administrative process involved in obtaining a non-honorable discharge, many members discharged for minor disciplinary reasons will actually be classified as an honorable discharge. Only serious offenses such as illicit drug possession, assault, and criminal misconduct will generally trigger non-honorable discharges. We find that the increase in discharges are driven by honorable discharges in the first year. Column 9 shows that honorable discharges increase by about 1.2 percentage point within the first year after the ED visit (9% increase), but there is no effect for non-honorable discharges. After two years, the effect for honorable discharges becomes slightly smaller, but we find a large and statistically significant increase in non-

honorable discharges of 0.8 percentage points (13% increase). It is not surprising that we find delayed effects for non-honorable discharges because it takes time for formal investigations to conclude, while honorable discharges can be implemented more expeditiously. While both categories include discipline-related discharges, the non-honorable discharges will contain more serious infractions while honorable discharges will be a mix of minor infractions and discharges unrelated to misconduct (e.g., for medical reasons, pregnancy, end of service term). Overall, the results in this section show large increases in discipline-related discharges from opioid use.

5.4 Heterogeneity Analyses

Next, we consider how the IV estimates vary across different subgroups. We consider subgroups based on demographic characteristics and pre-existing diagnoses of depression or anxiety.²⁷ Figure 2 shows the coefficient estimates and 95% confidence intervals from estimating the IV regression in equation (4) separately for each demographic subgroup.

The first two panels display the results for long-term opioid use (180 days supply) and doctor shopping. Receiving an opioid prescription in the ED predicts an increase in long-term opioid use and doctor shopping for almost all subgroups, although not all estimates are statistically significant. We find larger effects for White military members and those without a college education for doctor shopping, which is the strongest indicator of opioid misuse. These results are consistent with Case and Deaton (2015) which showed that White non-Hispanic individuals with less than a college education have been most impacted by the opioid crisis.

In the remaining panels of Figure 2, we examine heterogeneity in workforce outcomes. We focus on promotions, discipline flags, discipline discharges and non-honorable discharges for which we found the largest effects in the full sample. For promotions, although effects are roughly similar across subgroups, we find larger negative point estimates for military members who are White, have a below median AFQT score, and for those who have depression or anxiety. The most pronounced differences occur by race. White members experience a much larger negative effect of opioid use on the likelihood of a promotion, while the effect for non-White members is not statistically distinguishable from zero. When examining discipline flags,

²⁷ At baseline, 16% are receiving medication for depression or anxiety. Depression/Anxiety diagnoses are defined based on prescription fills in the year prior to the index ED visit for one of the following therapeutic classes: Benzodiazepines, Antidepressants, or Antipsychotic agents.

discipline discharges, and non-honorable discharges we do not find meaningful differential effects by race. Instead, the characteristics that most strongly predict a higher likelihood of disciplinary action or discharge are: younger than age 30, single, no college education, below median AFQT score, and having depression or anxiety. These results show that certain demographic factors predict a greater likelihood that initial opioid exposure results in negative performance and employment consequences.

5.5 Robustness Tests and Alternative Explanations

5.5.1 Sensitivity to Control Variables

In Appendix Table 3, we conduct a number of robustness checks to evaluate whether our main IV results hold up to varying the model specification. In Column 1, we present the results excluding all controls except for hospital-year-month and hospital-day of week fixed effects. In Column 2, we add diagnosis fixed effects. In Columns 3-6 we add basic demographic control variables, education, and military occupational variables. The results are generally robust to the set of controls included. The majority of coefficients do not change as control variables are added, suggesting coefficient stability, and some standard errors decrease. Almost all results remain statistically significant at the 5% level. Thus, the control variables may improve precision but do not materially impact our main coefficients, which is expected given random assignment.

5.5.2 Unobserved Injury Severity

Our instrumental variable analysis finds worse workforce outcomes for those who received an opioid in the ED. One alternative explanation for these findings may be that these individuals have a (unobservably) more serious or lasting injury that would independently lower job performance. If this was correlated with being assigned to a high propensity opioid prescriber, then this could bias our estimates upward in magnitude. The balance tests presented in Table 1 and Appendix Table 1 provide strong support that assignment to a high propensity prescriber is unrelated to patient characteristics. However, we additionally test this alternative hypothesis more directly by examining whether there are differences in health status within one year among patients who received an opioid prescription in the ED. While receiving an opioid prescription could have a direct causal impact on health status on its own, finding a null effect would suggest both that 1) there is no impact of the opioid itself on health status and 2) patient

assignment to physicians is uncorrelated with the severity of the medical condition, supporting the conditional independence assumption. We conduct this test using the Physical Capability Grading System score, which is derived from annual health assessments of Army members. In this assessment, a medical provider conducts a physical exam to assess a member's current health status and to identify any medical conditions. They also conduct a behavioral health screening. In Appendix Table 4, our IV estimates suggest that there is no effect of receiving an opioid prescription on the Z-scores for the physical and behavioral assessments in the year following the ED visit. Thus, members who received an opioid prescription are as physically capable as those who did not receive an opioid prescription, suggesting that the severity of the medical condition cannot explain the lower job performance outcomes we observe. These results are consistent with Table 7, which also showed that opioid receipt did not predict a higher likelihood of being discharged from the military for a medical reason.²⁸ Taken together, these results suggest that it is the receipt of an opioid prescription rather than the injury or medical condition itself that leads to negative workforce outcomes.

5.5.3 Exclusion Restriction

Another potential concern is that a physician's opioid prescribing propensity could be correlated with their propensity to provide other medical services during the ED visit which also impact downstream outcomes. To the extent there is a positive correlation in these propensities, additional medical services would most likely (weakly) improve health outcomes and work performance (which we did not observe in the previous section), hence biasing our estimates towards zero. To test for this, we construct a leave-out residualized instrument analogous to equation (2) for four measures of medical treatment intensity for the index ED visit: Berenson-Eggers Type of Service (BETOS) codes, BETOS and procedures, log work relative value units (wRVUs), and the ED prescribing propensity for anxiety or depression medication.²⁹ We

²⁸ While it may seem surprising that opioid use would not impact health status measures, we note that opioid abuse disorder diagnoses and non-fatal overdoses, which would trigger the largest changes in health care utilization from opioid misuse, are rare within the first few years after an ED prescription (Eichmeyer and Zhang (2022)). These null results are also in line with studies in the Danish setting that showed no increase in sick leave and disability claiming related to opioid prescribing (Laird and Nielsen (2017)).

²⁹ BETOS codes are clinically meaningful categories of HCPCS procedure and service codes. Specifically, we include the number of BETOS codes for procedures, diagnostic tests and imaging. We exclude Evaluation & Management (E&M) codes since nearly every visit includes one claim per patient, as well as durable medical equipment, and unclassified codes. We take the log of wRVUs given considerable skewness in this variable.

include the latter measure to account for non-opioid prescribing intensity that could have lasting effects beyond the acute condition.³⁰ We then include each predicted medical treatment intensity measure in the original 2SLS equation (4) as a control. Appendix Table 5 shows that our main results are generally robust to controlling for each measure of medical treatment intensity suggesting that receiving an opioid prescription predicts negative workforce outcomes independent of other medical services rendered during the ED visit.

5.5.4 Instrument Monotonicity

An assumption of our 2SLS strategy is that the relationship between physician prescribing propensities and the likelihood of receiving an opioid is monotonic. In other words, a patient that receives an opioid from a low propensity physician would also receive one from a high propensity physician. If the monotonicity assumption is violated, then we are unable to interpret our estimates as local average treatment effects (LATE). In Appendix Table 6, we conduct two tests used in prior studies to provide support for this assumption. In Column 1, following Eichmeyer and Zhang (2022), we estimate the first stage regression for age, gender, marital status, race, education, AFQT, and depression subsamples. This leads to 14 distinct subsamples in which we can observe whether there is a positive first-stage relationship. Across all subsamples we find a robust statistically significant positive relationship between the instrument (physician prescribing propensity) and the probability of receiving an opioid, consistent with the monotonicity assumption. In Column 2, we leave out each subsample in constructing the instrument and then run the first stage on the left-out subsample. Again, we find large and positive estimates across the subsamples.

5.5.5 Sample Attrition

Finally, we consider how attrition from the military affects our main estimates. About 18% of military members exit the sample within a year of the ED visit due to job separation. We explore how these individuals differ from the sample that does not attrit and we estimate the extent to which these discharges impact our main results. To do this, we re-estimate our results conditioning on members who stay in the military. Given that we find large effects of opioid receipt on job separations, those who leave the military are more likely to be misusing opioids

³⁰ For the young adult military population, anxiety and/or depression drugs are among the most commonly used chronic medications.

and have worse job performance. Thus, excluding attriters from our sample would likely understate the effects of receiving an opioid prescription on job performance outcomes. Specifically, we would understate the decrease in promotions and the increase in discipline flags.

Panel A of Appendix Table 7 displays our main results excluding attriters. Column 1 shows the full sample. Column 2 shows results excluding those who exit the military within 6 months after the ED visit and Column 3 excludes those who exit between 6 months and 1 year. Column 2 results are similar to the full sample. Thus, it appears that members who leave the military within six months after the ED visit do so for reasons unrelated to opioid use or adverse performance. This is not surprising given that it would take time for a person to transition from an initial opioid prescription to misuse and negative work behaviors. The results in Column 3 decrease slightly for opioid use and promotions, and there is a somewhat larger decrease for disciplinary flags. These results are consistent with our main results on job separations. They show that members with negative performance outcomes (especially a discipline flag) following an opioid prescription are most likely to be discharged. Thus, excluding these members would slightly understate the negative effects of opioid use on job performance outcomes.³¹ However, since these differences are small, our main results are relatively insensitive to whether or not we exclude attriters. Thus, the negative performance effects—reduction in promotions and increase in disciplinary actions—are not driven by the increase in discharges.

6. Conclusion

This paper adds to the nascent literature on the impact of opioid use on workforce outcomes. We focus our analysis on the U.S. military because of the unique ability to link medical records with detailed job performance outcomes. Relative to the existing literature, we provide the first evidence, to our knowledge, on the broader implications of opioid initiation on measures of job performance and capability. We investigate the relationship between opioid use and employment outcomes by employing a 2SLS identification strategy that relies on the quasi-

³¹ This conclusion is also supported by the summary statistics shown in Panel B of Appendix Table 7, which displays sample means for those who exit the military. Members who exited between 6-12 months after the ED visit have a much higher probability of long-term opioid use and doctor shopping relative to the full sample, even with the shorter follow-up period. These individuals are also more likely to have worse job performance outcomes. On the other hand, members who exited between 0-6 months are much less likely to have problematic opioid use (unsurprisingly since the follow-up period for these individuals is very short) and negative employment outcomes.

random assignment of patients to physicians in the ED and the physician's historic opioid prescribing propensity at the time of the patient's ED visit.

Our results have three main take-aways. First, we show that opioid initiation reduces job performance. We observe fewer promotions, increases in demotions, and fewer military members obtaining security clearances. Most of these negative outcomes emerge within one year after opioid initiation. Second, we show that opioid initiation leads to behavioral problems that increase the likelihood of disciplinary action and discharges. These findings suggest lower job productivity that can lead to increased recruitment and employee retention costs. Effects on disciplinary actions are concentrated among minor infractions, such as being late to work and unexcused absences. We do not, however, find increases in serious criminal activities or drug abuse. The latter result is consistent with the findings that the majority of opioid use is tied to legal prescriptions and that there is no observed effect of opioid use on the abuse of other drugs. While about half of the effect on discharges comes from increases in discipline-related separations, we also observe substantial increases in contract non-renewal or retirement. Third, while we observe negative workforce outcomes related to behavioral issues, we do not find reductions in physical capability following opioid initiation. However, our physical performance outcomes are conditional on testing, thus leaving open the possibility that some individuals with difficulties meeting physical expectations were able to avoid testing.

We use our results to perform a back-of-the-envelope calculation to estimate how opioid misuse and workforce outcomes would change at the population level if the physician opioid prescribing rate were reduced from the mean rate (22%) to the rate of the 25th percentile physician (15%) through policy levers.³² This extrapolation exercise suggests there would be a 4% decline in doctor shopping behavior and long term opioid use, a 0.3% increase in promotions and a 0.9% decrease in disciplinary flags among active-duty military members. Overall, discharges would decrease by 0.6% and discipline-related discharges would decrease by 1.8%.

³² This extrapolation scales our IV estimates by 0.07, the difference from moving between the mean rate (0.22) to the 25th percentile (0.15). Alternatively, we could extrapolate from an (out-of-sample) prescribing rate of zero to the mean rate (0.22), to represent how the introduction of the opioid epidemic may have affected workforce outcomes. In this case, we would find effects on opioid misuse and workforce outcomes that are about three-times as large. For example, moving from zero to the current prescribing rate would increase overall discharges by 2% and discipline-related discharges by 6%. As a point of reference, the overall U.S. employment rate fell by 6% from 1999 to 2018 (Abraham and Kearney (2020)).

In summary, these findings show that the opioid epidemic has had significant negative consequences for labor market productivity. They also suggest that opioid misuse has negatively impacted the military's readiness for future missions. Moreover, additional military resources have been dedicated to job performance reprimands that could have otherwise been used for productive workforce related efforts. Encouraging safer use of opioids and increasing substance abuse treatment has the potential to reduce job loss and increase productivity. Although employer spending on substance abuse treatment has increased in recent years—large employer health plans spent \$2.6 billion on treatment for opioid addiction and overdoses in 2016, up from \$828 million in 2010 (KFF, 2018)—further expansions may be warranted. Identifying those in need of opioid treatment in the workplace, however, is challenging. We show that the behavioral consequences of opioid misuse are more likely to trigger disciplinary actions and separations than drug and medical screenings.

Our paper has several limitations. First, we estimate the effects of opioid initiation for active duty military members receiving care on U.S. military bases. This may not generalize to military members serving in combat overseas or to veterans, though previous work has shown similar opioid use patterns for veterans (Barnett et al. (2019); Zhang (2021)). However, our sample more closely resembles civilian workers in physically demanding jobs relative to other military populations, thus broadening its applicability. Second, it is possible that our results reflect lower bound estimates of negative workforce outcomes since negative work performance requires a formal administrative process to record disciplinary actions. Some commanding officers may not pursue this process for minor infractions, thus, we may not capture all behavioral problems. Finally, our 2SLS strategy relies on the assumption that our instrument does not operate through channels other than opioid prescribing in the ED. In particular, a physician who is a high-intensity opioid prescriber may also provide other types of care to the patient more (or less) intensively. Although we cannot control for unobserved components of patient care, we find that opioid prescribing independently predicts long-term opioid use and negative workforce outcomes when controlling for observable measures of medical care intensity.

References


Abraham, K.G. and Kearney, M.S. (2020). Explaining the decline in the US employment-to-population ratio: A review of the evidence. *Journal of Economic Literature*, 58(3), pp. 585-643.

- AJMC Perspectives (2020). Economic burden of illness in opioid use disorder (OUD) and medication-assisted treatments.
- Aliprantis, D., Fee, K., and Schweitzer, M. (2019). Opioids and the labor market. Working Paper 18-07R, Federal Reserve Bank of Cleveland.
- Alpert, A. E., Dykstra, S. E., & Jacobson, M. (2020). How do prescription drug monitoring programs reduce opioid prescribing? The role of hassle costs versus information. NBER Working Paper 27584.
- Alpert, A., Powell, D., & Pacula, R. L. (2018). Supply-side drug policy in the presence of substitutes: Evidence from the introduction of abuse-deterrent opioids. *American Economic Journal: Economic Policy*, 10(4), 1-35.
- Barnett, M.L., Olenski, A.R., and Jena, A.B. (2017). Opioid-prescribing patterns of emergency physicians and risk of long-term use. *New England Journal of Medicine*, 376(7), pp. 663–673.
- Barnett, M.L., et al. (2019). Emergency physician opioid prescribing and risk of long-term use in the veterans health administration: an observational analysis. *Journal of General Internal Medicine*, 34(8), pp. 1522-1529.
- Beheshti, D. (2022). The impact of opioids on the labor market: Evidence from drug rescheduling. *Journal of Human Resources*, 0320-10762R1.
- Buchmueller, T.C. and Carey, C. (2018). The effect of prescription drug monitoring programs on opioid utilization in Medicare. *American Economic Journal: Economic Policy*, 10(1), 77-112.
- Butikofer, A. and Skira, M.M. (2018). Missing work is a pain: The effect of cox-2 inhibitors on sickness absence and disability pension receipt. *Journal of Human Resources*, 53 (1), 71–122.
- Case, A. and Deaton, A. (2015). Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proceedings of the National Academy of Sciences*, 112(49), pp.15078-15083.
- Centers for Disease Control and Prevention (CDC). (2019). Opioids in the workplace: data. <https://www.cdc.gov/niosh/topics/opioids/data.html>. February 27, 2022.
- Centers for Disease Control and Prevention (CDC). (2020). Acute pain. <https://www.cdc.gov/acute-pain/about/index.html> February 27, 2022.
- Centers for Disease Control and Prevention (CDC). (2021). Opioid overdose: Understanding the epidemic. <https://www.cdc.gov/drugoverdose/epidemic/index.html>. February 27, 2022.
- Cesur R., Sabia J.J., and Bradford, W.D. (2019). Did the war on terror ignite an opioid epidemic? NBER Working Paper 26264.
- Chan, D. (2016). Teamwork and moral hazard: Evidence from the emergency department. *Journal of Political Economy*, 124(3), pp. 734-770.
- Chang, C. and Obermeyer, Z. (2020). Association of clinical characteristics with variation in emergency physician preferences for patients. *JAMA network open*, 3(1), e1919607-e1919607.

- Cho, D., Garcia, D., Montes, J., & Weingarden, A. (2021). Labor Market Effects of the Oxycodone-Heroin Epidemic. Finance and Economics Discussion Series 2021-025. Washington: Board of Governors of the Federal Reserve System
- Cone, E.J., Welch, P., Mitchell, J.M., and Paul, B.D. (1991). Forensic drug testing for opiates: I. Detection of 6-acetylmorphine in urine as an indicator of recent heroin exposure; drug and assay considerations and detection times. *Journal of Analytical Toxicology*, 15(1), pp.1-7.
- Currie, J. and Madrian, B. C. (1999). Health, health insurance and the labor market. *Handbook of Labor Economics*, 3, pp. 3309-3416.
- Dave, D., Deza M., and Horn, B. (2021). Prescription drug monitoring programs, opioid abuse, and crime. *Southern Economic Journal*, 87(3), pp. 808-48.
- Deiana, C. and Giua, L. (2021). The intended and unintended effects of opioid policies on prescription opioids and crime. *The BE Journal of Economic Analysis & Policy*, 21(2), 751-92.
- Eichmeyer, S. and Zhang, J. (2023). Primary care provider's influence on opioid use and its adverse consequences. *Journal of Public Economics*, forthcoming.
- Eichmeyer, S. and Zhang, J. (2022). Pathways into opioid dependence: Evidence from practice variation in emergency departments. *American Economic Journal: Applied Economics*, 14(4): 271-300.
- Garthwaite, C. (2012). The economic benefits of pharmaceutical innovations: The case of cox-2 inhibitors. *American Economic Journal: Applied Economics*, 4(3), pp. 116–37.
- Harris, M.C, Kessler, L.M., Murray, M.N., and Glenn, B. (2020). Prescription opioid and labor market pains: The effect of schedule II opioids on labor force participation and unemployment. *Journal of Human Resources*, 55(4), pp. 1319-1364.
- Hernandez, I., He, M., Brooks, M. M., & Zhang, Y. (2018). Exposure-response association between concurrent opioid and benzodiazepine use and risk of opioid-related overdose in Medicare Part D beneficiaries. *JAMA network open*, 1(2), e180919-e180919.
- Hollingsworth, A., Ruhm, C., and Simon, K. (2017). Macroeconomic conditions and opioid abuse. *Journal of Health Economics*, 56, pp. 222-233.
- Kaiser Family Foundation. (2018). A look at how the opioid crisis has affected people with employer coverage.
- Krueger, A. B. (2017). Where have all the workers gone? An inquiry into the decline of the U.S. labor force participation rate. *Brookings Papers on Economic Activity*, 2, pp. 1–87.
- Laird, J. and Nielsen, T. (2017). The effects of physician prescribing behaviors on prescription drug use and labor supply: Evidence from movers in Denmark. Mimeo, Harvard University.
- Lin, L.A., et al. (2019). Changing trends in opioid overdose deaths and prescription opioid receipt among veterans. *American Journal of Preventive Medicine*, 57 (1), pp. 106-110.
- Maclean, J.C., Mallatt, J., Ruhm, C.J., and Simon, S. (2021). Economic studies on the opioid crisis: A review. NBER Working Paper 28067.


- Mallatt, J. (2018). The effect of prescription drug monitoring programs on opioid prescriptions and heroin crime rates. SSRN Working Paper.
- Mallatt, J. (2020). Policy-induced substitution to illicit drugs and implications for law enforcement activity. SSRN Working Paper.
- Mayo Clinic (2020). Exercise and opioids: What to know before you work out. <https://www.mayoclinic.org/healthy-lifestyle/consumer-health/in-depth/opioids-and-exercise/art-20396866>. February 27, 2022.
- Mazereeuw, G., Sullivan, M.D., and Juurlink, D.N. (2018). Depression in chronic pain: Might opioids be responsible? *Pain*, 159(11), pp. 2142-2145.
- Meinhofer, A. (2016). The war on drugs: Estimating the effect of prescription drug supply-side interventions. SSRN Working Paper.
- Meyer, C. (2019). Opioids in the workplace: Impairment, illicit substances, and insider threat, *Security Management*.
- National Institute of Drug Abuse (2020). <https://www.drugabuse.gov/drug-topics/health-consequences-drug-misuse/mental-health-effects>. February 27, 2022.
- Park, S. and Powell, D. (2021). Is the rise in illicit opioids affecting labor supply and disability claiming rates? *Journal of Health Economics*, 76, 102430.
- Peters, Z.J., et al. (2019). Surveillance snapshot: Trends in opioid prescription fills among U.S. military service members during fiscal years 2007–2017, *Health.mil*.
- Pierce, J. and Schott, P. (2020). Trade liberalization and mortality: Evidence from U.S. counties. *American Economic Review: Insights*, 2(1), pp. 47-64.
- Powell, D. (2021). The labor supply consequences of the opioid crisis. Rand Working Paper.
- Powell, D., Pacula, R. L., & Jacobson, M. (2018). Do medical marijuana laws reduce addictions and deaths related to pain killers?. *Journal of Health Economics*, 58, 29-42.
- Richards G.C., et al. (2018). Effects of long-term opioid analgesics on cognitive performance and plasma cytokine concentrations in patients with chronic low back pain: a cross-sectional pilot study. *Pain Reports* 3(4), p. E669.
- Rockeman, O. (2021). The mystery of the missing workers, explained, *Bloomberg*.
- Smith, K. (2021). <https://www.psychom.net/depression.central.opioid.abuse.html>.
- Thingholm, P. (2020). Physician heterogeneity and patient outcomes. PhD Dissertation, Aarhus University.
- Venkataramani A.S., Bair E.F., O'Brien R.L., and Tsai, A.C. (2020). Association between automotive assembly plant closures and opioid overdose mortality in the United States: A difference-in-differences analysis. *JAMA Internal Medicine*, 180(2), pp. 254-62.
- Winkelman T.N., Chang V.W., and Binswanger, I.A. (2018). Health, polysubstance use, and criminal justice involvement among adults with varying levels of opioid use. *JAMA Network Open*, 1(3), e180558.

Figure 1: Distribution of Physician Opioid Prescribing Rate



Notes: Figure displays the distribution of physicians’ mean opioid prescribing rate for each year (labeled as “raw”). The residualized prescribing rate for each physician-year is shaded which reflects idiosyncratic factors affecting prescribing decisions that are unrelated to hospital, time, diagnosis, and patient sociodemographic characteristics.

Figure 2: Heterogeneity of IV Results



Notes: Figure displays subsample analysis of the instrumental variable results using the prescribing rate instrument for six main outcomes. Non-Honorable Discharge (2 Yr) includes all non-honorable discharges within 2 years of the ED visit (“General”, “Other than Honorable,” “Bad Conduct”, and “Dishonorable”).

Table 1: Descriptive Statistics and Balance Test

	Full Sample Mean	Prescribing Intensity			High Prescribing Intensity		
		Coefficient	Std. Error	p-value	Coefficient	Std. Error	p-value
	(1)	(2)	(3)	(4)	(5)	(6)	(7)
Age	26.17	-0.116	0.448	0.795	-0.034	0.085	0.688
Female	0.21	-0.030	0.025	0.235	0.002	0.005	0.747
Race - White	0.65	0.007	0.031	0.808	0.001	0.006	0.813
Rank - Junior	0.62	-0.003	0.033	0.932	-0.001	0.006	0.823
College	0.05	-0.006	0.009	0.526	-0.002	0.002	0.284
Marital Status - Married	0.53	0.003	0.023	0.895	0.000	0.004	0.937
Military Tenure (Yrs)	5.44	-0.011	0.357	0.976	0.008	0.069	0.908
AFQT	59.82	-0.047	1.628	0.977	0.191	0.283	0.500
Observations	1,447,758	1,447,758	1,447,758	1,447,758	647,123	647,123	647,123

Notes: Means in Column 1 are calculated based on the patient’s characteristics at the time of the ED visit. Coefficients, standard errors, and p-values stem from a regression with the demographic characteristic as the dependent variable and the residualized instrument (continuous Prescribing Intensity instrument in Columns 2-4 and binary indicator for High Prescribing Intensity in Columns 5-7) as the independent variable. Each cell represents results from a separate regression and standard errors are clustered by physician.

Table 2: Probability of Opioid Prescription in ED and Long-Term Opioid Use

	First Stage	2SLS - Opioid Use Within 1 Year		
	Prescription in ED (1)	Number of Prescriptions (2)	Doctor Shopping (3)	Long Term Use (180 Days Supply) (4)
<i>Panel A: Prescribing Rate Instrument</i>				
Prescribing Intensity	0.8644*** (0.009)			
Prescription in ED		0.0384** (0.0165)	0.0018** (0.0008)	0.0018** (0.0008)
<i>Panel B: High vs. Low Instrument</i>				
High Prescribing Intensity	0.1847*** (0.0033)			
Prescription in ED		0.0613*** (0.0152)	0.0018** (0.0007)	0.0022*** (0.0007)
Full Sample Mean	0.22	0.3603	0.0030	0.0029
Total Observations	1,447,758	1,447,758	1,447,758	1,447,758

Notes: This table presents coefficients obtained from the first stage (Column 1) and the second stage (Columns 2-4) of the instrumental variable regressions on the impact of an opioid prescription on long-term opioid use after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Doctor shopping is defined as having 7 or more different prescribers in 1 year. Number of prescriptions and long term use exclude the first prescription filled within 7 days of ED visit. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Table 3: IV Results for Probability of Drug Test Failure

	Drug Test Failure Involving:						
	Drug Test Failure	Opioid with Prescription	Opioid without Prescription	Heroin	Marijuana	Benzos	Other Drugs
	(1)	(2)	(3)	(4)	(5)	(6)	(7)
<i>Panel A: Prescribing Rate Instrument</i>							
Prescription in ED	0.0100*** (0.0033)	0.0076*** (0.0018)	0.0003 (0.0010)	-0.0005** (0.0002)	0.0001 (0.0018)	0.0000 (0.0007)	0.001 (0.0016)
<i>Panel B: High vs. Low Instrument</i>							
Prescription in ED	0.0104*** (0.0028)	0.0088*** (0.0014)	0.0009 (0.0009)	-0.0003 (0.0002)	0.0002 (0.0017)	-0.0003 (0.0006)	-0.0003 (0.0015)
Full Sample Mean	0.029	0.0068	0.0032	0.00024	0.012	0.0016	0.0086
Total Observations	1,045,277	1,045,277	1,045,277	1,045,277	1,045,277	1,045,277	1,045,277

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on the probability of drug test failure in the year after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Table 4: IV Results for Probability of Promotion and Demotion

	Promotion (1 Yr)	Promotion (2 Yr)	Demotion (1 Yr)	Demotion (2 Yr)
	(1)	(2)	(3)	(4)
<i>Panel A: Prescribing Rate Instrument</i>				
Prescription in ED	-0.0135** (0.0057)	-0.0120* (0.0062)	0.0005 (0.0027)	0.0019 (0.0031)
<i>Panel B: High vs. Low Instrument</i>				
Prescription in ED	-0.0137** (0.0054)	-0.0112* (0.0059)	0.0008 (0.0026)	0.0027 (0.0030)
Full Sample Mean	0.28	0.39	0.035	0.047
Total Observations	1,447,750	1,447,750	1,447,750	1,447,750

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on the probability of promotion and demotion within 1 and 2 years after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Table 5: IV Results for Army Physical Fitness Test (APFT) - Army Only

	Take an APFT (1)	Pass an APFT (2)	Z-Score APFT (3)
<i>Panel A: Prescribing Rate Instrument</i>			
Prescription in ED	-0.0092 (0.0089)	-0.0068 (0.0062)	-0.0267 (0.0227)
<i>Panel B: High vs. Low Instrument</i>			
Prescription in ED	-0.0149* (0.0083)	-0.0048 (0.0057)	-0.0353 (0.0223)
Full Sample Mean	0.57	0.93	-
Total Observations	718,973	410,033	409,946

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on Army Physical Fitness test outcomes in the year after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Table 6: IV Results for Disciplinary Actions - Army Only

	Discipline Flag (1)	Criminal Investigation Flag (2)	Drug or Alcohol Flag (3)
<u>Within 1 Year Following ED Visit:</u>			
<i>Panel A: Prescribing Rate Instrument</i>			
Prescription in ED	0.0108** (0.0052)	0.0008 (0.0011)	0.0005 (0.0018)
<i>Panel B: High vs. Low Instrument</i>			
Prescription in ED	0.0110** (0.0051)	0.0012 (0.0012)	0.0005 (0.0015)
Full Sample Mean	0.087	0.0058	0.0082
<u>Within 2 Years Following ED Visit:</u>			
<i>Panel A: Prescribing Rate Instrument</i>			
Prescription in ED	0.0084 (0.0052)	0.0017 (0.0015)	0.0012 (0.0021)
<i>Panel B: High vs. Low Instrument</i>			
Prescription in ED	0.0086 (0.0056)	0.0019 (0.0016)	0.0008 (0.0019)
Full Sample Mean	0.12	0.0096	0.012
Total Observations	718,973	718,973	718,973

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on the probability of an Army “Suspension of Favorable Personnel Actions (SFPA) Flag” within 1 and 2 years after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Table 7: IV Results for Military Discharges

	Reason for Discharge:						Character of Separation:				
	Any Discharge	Non-Renewal of Contract or Retirement		Failure to Meet Weight or Body Fat Standards		Substance Abuse	Other Non-Discipline	Medical	Uncharacterized	Honorable	Non-Honorable
		(1)	(2)	(3)	(4)						
<u>Within 1 Year Following ED Visit:</u>											
<i>Panel A: Prescribing Rate Instrument</i>											
Prescription in ED	0.0159** (0.0065)	0.0113*** (0.0036)	0.0067*** (0.0022)	0.0010 (0.0011)	-0.0008 (0.0017)	0.0007 (0.0019)	-0.0011 (0.0028)	-0.0019 (0.0024)	0.0117** (0.0055)	0.0052 (0.0033)	
<i>Panel B: High vs. Low Instrument</i>											
Prescription in ED	0.0126** (0.0055)	0.0091*** (0.0034)	0.0066*** (0.0020)	0.0009 (0.0010)	-0.0024 (0.0016)	0.0003 (0.0017)	-0.0006 (0.0025)	-0.0013 (0.0021)	0.0086* (0.0049)	0.0049* (0.0029)	
Full Sample Mean	0.18	0.068	0.026	0.0045	0.014	0.017	0.027	0.022	0.13	0.043	
<u>Within 2 Years Following ED Visit:</u>											
<i>Panel A: Prescribing Rate Instrument</i>											
Prescription in ED	0.0126* (0.0072)	0.0071 (0.0047)	0.0069** (0.0028)	0.0005 (0.0014)	-0.001 (0.0021)	0.0022 (0.0027)	-0.0016 (0.0036)	-0.0015 (0.0026)	0.0058 (0.0066)	0.0084** (0.0038)	
<i>Panel B: High vs. Low Instrument</i>											
Prescription in ED	0.0158** (0.0064)	0.0075* (0.0045)	0.0080*** (0.0026)	0.0009 (0.0013)	(0.0019) (0.0020)	0.0013 (0.0023)	0.0008 (0.0032)	(0.0006) (0.0024)	0.0084 (0.0059)	0.0092*** (0.0034)	
Full Sample Mean	0.32	0.14	0.042	0.0077	0.02	0.031	0.054	0.029	0.25	0.063	
Total Observations	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	1,447,758	

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on the probability of military discharge within 1 and 2 years after the ED visit, reason for discharge and “character of separation.” Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Honorable + non-honorable discharge do not add up to any discharge because there are some discharges that are unclassified. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

FOR ONLINE PUBLICATION

Appendix

Appendix Table 1: Joint Balance Test of Patient Characteristics and Physician Opioid Prescribing Intensity

	(1)	(2)	(3)
	ED Prescription	Residualized Instrument	Residualized Instrument (incl. time, hospital, diagnosis)
Age	0.0028*** (0.0002)	0.0000 (0.0001)	0.0000 (0.0001)
Race - White	0.0350*** (0.0014)	0.0000 (0.0007)	0.0000 (0.0007)
Female	-0.0434*** (0.0017)	-0.0009 (0.0007)	-0.0008 (0.0007)
Rank - Junior	-0.0218*** (0.0017)	-0.0002 (0.0007)	-0.0002 (0.0007)
Education - College Degree	-0.0293*** (0.0020)	-0.0003 (0.0007)	-0.0003 (0.0007)
Marital Status - Married	0.0117*** (0.0011)	0.0001 (0.0004)	0.0001 (0.0004)
Military Tenure (Yrs)	0.0012*** (0.0002)	0.0000 (0.0001)	0.0000 (0.0001)
Armed Forces Qualificaiton Test Percen	-0.0005*** (0.0000)	0.0000 (0.0000)	0.0000 (0.0000)
Observations	1,447,758	1,447,758	1,447,758
F-Statistic	611.62	0.27	0.28

Notes: Column 1 shows a regression of a binary indicator for receiving an opioid prescription on patient characteristics. Column 2 shows the balance test where we regress the continuous physician prescribing rate instrument on patient characteristics. Column 3 is the same as Column 2 except that we construct the instrument using only hospital-month-year, hospital-day of week, and diagnosis fixed effects (i.e., sociodemographic controls are excluded from the instrument). Consistent with random assignment, patient characteristics do not predict the physician opioid prescribing intensity. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 2: IV Results for Probability of Obtaining a Security Clearance – Army Only

	Within 1 Year following ED Visit		Within 2 Years following ED Visit	
	Denied a Security Clearance (1)	Security Clearance Revoked (2)	Denied a Security Clearance (3)	Security Clearance Revoked (4)
<i>Panel A: Prescribing Rate Instrument</i>				
Prescription in ED	0.0047** (0.0021)	0.001 (0.0021)	0.0041* (0.0024)	-0.0011 (0.0026)
<i>Panel B: High vs. Low Instrument</i>				
Prescription in ED	0.0045** (0.0021)	0.0001 (0.0020)	0.0040* (0.0024)	-0.0005 (0.0023)
Full Sample Mean	0.017	0.012	0.021	0.018
Total Observations	718,973	718,973	718,973	718,973

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on the probability of security clearance outcomes within 1 and 2 years after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. Full sample mean and total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 3: Sensitivity to Control Variable Selection

<i>Specifications: Prescribing Rate Instrument Results</i>	(1)	(2)	(3)	(4)	(5)	(6)
Prescriptions in 1 Year	0.0323* (0.0165)	0.0357** (0.0168)	0.0371** (0.0166)	0.0370** (0.0166)	0.0371** (0.0166)	0.0384** (0.0165)
7 Providers in 1 Year	0.0017** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)
180 Days of Supply in 1 Year	0.0017** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)
Promotion in 1 Year	-0.0100 (0.0090)	-0.0155* (0.0079)	-0.0142** (0.0064)	-0.0143** (0.0064)	-0.0102* (0.0058)	-0.0135** (0.0057)
Disciplinary Flag in 1 Year (Army Enlisted Only)	0.0156** (0.0063)	0.0122** (0.0054)	0.0109** (0.0053)	0.0114** (0.0053)	0.0107** (0.0052)	0.0108** (0.0052)
<i>Fixed Effects</i>						
Hospital-Year-Month	Yes	Yes	Yes	Yes	Yes	Yes
Hospital-Day of Week	Yes	Yes	Yes	Yes	Yes	Yes
Diagnosis	No	Yes	Yes	Yes	Yes	Yes
Military Occupational Specialty (MOS)	No	No	No	No	No	Yes
<i>Controls</i>						
Age	No	No	Yes	Yes	Yes	Yes
Race - White	No	No	Yes	Yes	Yes	Yes
Female	No	No	Yes	Yes	Yes	Yes
Marital Status - Married	No	No	Yes	Yes	Yes	Yes
College	No	No	No	Yes	Yes	Yes
Rank - Junior	No	No	No	No	Yes	Yes
Military Tenure (Yrs)	No	No	No	No	Yes	Yes
AFQT Qualification Test Percentile	No	No	No	No	Yes	Yes

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions (using the residualized continuous prescribing rate as the instrument). Each cell represents a separate regression. The outcome variables are number of prescriptions, doctor shopping, long term opioid use, promotions and disciplinary action in the year after the ED visit. The first column includes only hospital-year-month, hospital-day of week fixed effects and each subsequent column adds control variables and/or fixed effects. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 4: IV Results for Physical Capability – Army Only

	(1)	(2)
	Physical Z-Score	Psych Z-Score
<i>Panel A: Prescribing Rate Instrument</i>		
Prescription in ED	-0.0083 (0.0208)	-0.0087 (0.0220)
<i>Panel B: High vs. Low Instrument</i>		
Prescription in ED	-0.0083 (0.0185)	-0.0088 (0.0195)
Total Observations	718,973	718,973

Notes: This table presents coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on physical and psychiatric capability (as measured by the Physical Capability Grading System score) within the year after the ED visit. Panel A uses the residualized continuous prescribing rate as an instrument while Panel B uses the binary version of the same variable (equal to one if the residualized prescribing rate is in the top quartile and zero if it is in the bottom quartile) for the endogenous variable of whether a patient filled an opioid prescription within 7 days of the ED visit. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. The raw physical and psych scale ranges from 1 to 5 with lower scores implying higher capability. Total observations are for the full sample used in Panel A. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 5: Controlling for Intensity of Medical Treatment during ED visit

	Main Results	Controlling for BETOS Number of Tests and Images Propensity	Controlling for BETOS Number of Tests, Images, and Procedures Propensity	Controlling for Log of Work RVU Propensity	Controlling for Medication Propensity
	(1)	(2)	(3)	(4)	(5)
Prescriptions in 1 Year	0.0384** (0.0165)	0.0368** (0.0162)	0.0324** (0.0163)	0.0323* (0.0169)	0.0363** (0.0165)
7 Providers in 1 Year	0.0018** (0.0008)	0.0017** (0.0008)	0.0017** (0.0008)	0.0015* (0.0008)	0.0018** (0.0008)
180 Days of Supply in 1 Year	0.0018** (0.0008)	0.0018** (0.0008)	0.0017** (0.0008)	0.0018** (0.0008)	0.0018** (0.0008)
Promotion in 1 Year	-0.0135** (0.0057)	-0.0128** (0.0056)	-0.0090* (0.0054)	-0.0087 (0.0055)	-0.0116** (0.0056)
Discipline Flag in 1 Year (Army Only)	0.0108** (0.0052)	0.0103** (0.0052)	0.0081 (0.0053)	0.0102* (0.0052)	0.0104** (0.0052)

Notes: This table presents in the first column the baseline coefficients obtained from the second stage of the instrumental variable regressions on our outcomes of interest: long-term opioid use, promotions and disciplinary flags. Column 1 displays the results from the previous baseline results. Columns 2 and 3 present 2SLS results that include the residualized BETOS control variable that measures the intensity of treatment with the performed number of tests and images during the ED visit and the number of performed tests, images, and procedures during the ED visit, respectively. Column 4 displays results controlling for the residualized log work RVU measure. The last column presents results from controlling for the residualized ED prescribing propensity for anxiety/depression medication. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 6: Monotonicity Tests

	Subsample First Stage (1)	Reverse-Sample Instrument (2)
Age < 30	0.8430*** (0.0094)	0.4647*** (0.0146)
Age 30+	0.9306*** (0.0123)	0.8557*** (0.0192)
White	0.8735*** (0.0091)	0.6500*** (0.0177)
Non-White	0.8497*** (0.0125)	0.7237*** (0.0212)
Female	0.7774*** (0.0197)	0.7067*** (0.0256)
Male	0.8851*** (0.0084)	0.5063*** (0.0167)
Single	0.8178*** (0.0113)	0.6556*** (0.0180)
Married	0.9060*** (0.0091)	0.7498*** (0.0193)
College	0.8258*** (0.0252)	0.8137*** -0.0351
Less than College	0.8662*** (0.0085)	0.1578*** -0.0088
AFQT Below Median Score	0.8813*** (0.0088)	0.7307*** (0.0160)
AFQT Above Median Score	0.8457*** (0.0109)	0.6882*** (0.0171)
Depression	0.8785*** (0.0212)	0.8024*** (0.0271)
No Depression	0.8601*** (0.0095)	0.3813*** (0.0175)
Full Sample	0.8644*** (0.0086)	-

Notes: This table displays results from two versions of the monotonicity test. In Column 1 we estimate the first stage by regressing the probability of receiving an opioid within 7 days of the ED visit on the continuous residualized prescribing rate instrument separately for each of the socio-demographic subsamples. In column 2 we leave out each respective subsample in constructing the residualized prescribing rate instrument and then run the first stage regression on the left-out subsample. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively.

Appendix Table 7: Attrition Tests

<i>Panel A: IV Results</i>	Full Sample	Excl. Attrition <6 months	Excl. Attrition 6-12 months
	(1)	(2)	(3)
Prescriptions in 1 year	0.0384** (0.0165)	0.0447** (0.0177)	0.0365** (0.0174)
7 Providers in 1 year	0.0018** (0.0008)	0.0019** (0.0008)	0.0018** (0.0008)
180 days of supply in 1 year	0.0018** (0.0008)	0.0021** (0.0009)	0.0019** (0.0008)
Promotion in 1 year	-0.0135** (0.0057)	-0.0136** (0.0059)	-0.0105* (0.0058)
Discipline flag in 1 year (Army only)	0.0108** (0.0052)	0.0120** (0.0055)	0.0049 (0.0051)
<i>Panel B: Means for Members who Attrit</i>	Full Sample Mean	Attrit <6 months	Attrit 6-12 months
	(1)	(2)	(3)
Prescriptions in 1 year	0.3603	0.0233	0.2796
7 Providers in 1 year	0.0030	0.0007	0.0040
180 days of supply in 1 year	0.0028	0.0007	0.0044
Promotion in 1 year	0.2796	0.0233	0.1044
Discipline flag in 1 year (Army only)	0.0867	0.0792	0.1901
Total Observations	1,447,758	121,526	136,944

Notes: This table presents in Panel A coefficients obtained from the second stage of the instrumental variable regressions on the impact of an opioid prescription on long-term opioid use, promotions and disciplinary action in the year after the ED visit. Column 1 displays the results from the previous baseline results for the full sample, while Columns 2 and 3 exclude those who exit the military within 6 months and 6 months to 1 year after the ED visit, respectively. Panel A uses the residualized continuous prescribing rate as an instrument. All regressions include the full set of fixed effects (hospital-month-year, hospital-day of week, diagnosis, and military service-by-occupational specialty) and the sociodemographic control variables. *, **, *** display statistical significance at the 10%, 5%, and 1% level respectively. Panel B displays mean outcomes for the full sample, and those that attrit within 6 months and 6 months to 1 year, respectively.